

ESSENTIAL ARTICLES OF PM&R

GAIT, ASSISTIVE DEVICES, PROSTHESES, ORTHOSES, and AMPUTATION-RELATED ISSUES

Authors:

Mark Hirsch, PhD

Vu Nguyen, MD

Gait

The effects of parallel bars, body weight support and speed on the modulation of the locomotor pattern of spastic paretic gait. A preliminary communication.

Visitin M, Barbeau H. Paraplegia 1994;32:540-553.

Gait variability and fall risk in community-living older adults: A 1-year prospective study.

Hausdorff JM, Rios DA, Edelberg HK. Archives of Physical Medicine and Rehabilitation 2001;82(8):1050-1056.

Hemiplegic gait – Analysis of temporal variables.

Brandstater ME, Debruin H, Gowland C, Clark BM. Archives of Physical Medicine and Rehabilitation 1983;64(12):583-587.

Relationships between spasticity, strength, gait, and the GMFM-66 in persons with spastic diplegia cerebral palsy.

Ross SA, Engsberg JR. Archives of Physical Medicine and Rehabilitation 2007;88(9):1114-1120.

Walking speed as a basis for normal and abnormal gait measurements.

Andriacchi TP, Ogle JA, Galante JO. Journal of Biomechanics 1977;10(4):261-268.

An evaluation of patient perceptions to the value of the gait laboratory as part of the rehabilitation of primary lower limb amputees.

Cole MJ, Durham S, Ewins D. Prosthetics and Orthotics International 2008;32(1): 12-22.

Gait efficiency using the C-Leg.

Orendurff MS, et al. Journal of Rehabilitation Research and Development 2006;43(2):239-46.

Kinematic and kinetic comparisons of transfemoral amputee gait using C-Leg and Mauch SNS prosthetic knees.

Segal AD, et al. Journal of Rehabilitation Research and Development 2006;43(7):857-70.

Differences in gait characteristics between persons with bilateral transtibial amputations, due to peripheral vascular disease and trauma, and able-bodied ambulators.

Su PF, et al. Archives of Physical Medicine and Rehabilitation 2008;89(7):1386-94.

Assistive Devices and Mobility

Mobile arm supports: Evidence-based benefits and criteria for use.

Atkins MS, et al. The journal of spinal cord medicine 2008;31(4):388-93.

Participation in community-based activities of daily living: Comparison of a pushrim-activated, power-assisted wheelchair and a power wheelchair.

Giesbrecht EM, et al. Disability and Rehabilitation Assistive Technology 2009;4(3):198-207.

Evaluation of a stair-climbing power wheelchair in 25 people with tetraplegia.

Laffont I, et al. Archives of Physical Medicine and Rehabilitation 2008;89(10):1958-64.

How many people would benefit from a smart wheelchair?

Simpson RC, LoPresti EF, Cooper RA. Journal of Rehabilitation Research and Development 2008;45(1):53-71.

Characterization of power wheelchair use in the home and community.

Sonenblum SE, et al. Archives of Physical Medicine and Rehabilitation 2008;89(3):486-91.

Orthoses

Orthotic and prosthetic prescriptions for today and tomorrow.

Phys Med Rehabil Clin N Am. Nov 2007;18(4):785-858. *This chapter is a nice overview of orthotic and prosthetic equipment.*

Energy expenditure in stroke subjects walking with a carbon composite ankle foot orthosis.

Danielsson A, Sunnerhagen KS. Journal of Rehabilitation Medicine 2004;36(4):165-8.

Effectiveness of upper and lower limb casting and orthoses in children with cerebral palsy: An overview of review articles.

Autti-Ramo I, et al. American Journal of Physical Medicine & Rehabilitation 2006;85(1): 89-103.

Effect of ankle-Foot orthoses on walking efficiency and gait in children with cerebral palsy.

Brehm MA, Harlaar J, Schwartz M. Journal of Rehabilitation Medicine 2008;40(7):529-34.

Effects of an anterior ankle-foot orthosis on postural stability in stroke patients with hemiplegia.

Chen CK, et al. American Journal of Physical Medicine & Rehabilitation 2008;87(10):815-20.

Indications for orthoses to improve gait in children with cerebral palsy.

Davids JR, Rowan F, Davis BR. The Journal of the American Academy of Orthopedic Surgeons 2007;15(3):178-88.

Orthotic management of the neuropathic limb.

Elftman NW. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):115-57.

Efficacy of ankle-foot orthoses on gait of children with cerebral palsy: Systematic review of literature.

Figueiredo EM, et al. Pediatric Physical Therapy 2008;20(3):207-23.

Effects of rigid and dynamic ankle-foot orthoses on normal gait.

Guillebastre BP, Calmels, Rougier P. Foot & Ankle International 2009;30(1):51-6.

Biomechanical effect of electromechanical knee-ankle-foot-orthosis on knee joint control in patients with poliomyelitis.

Hwang S, et al. Medical & Biological Engineering & Computing 2008;44(6):541-9.

Gait changes over time in stance control orthosis users.

Irby SE, Bernhardt KA, Kaufman KR. Prosthetics and Orthotics International 2007;31(4):353-61.

Shoe modification and the use of orthoses in the treatment of foot and ankle pathology.

Janisse DJ, Janisse E. Journal of the American Academy of Orthopaedic Surgeons 2008;16(3):152-8.

Advances in orthotics and bracing.

Logue JD. Foot and Ankle Clinics 2007;12(2):215-32.

Foot orthoses in rehabilitation-what's new.

Nawoczenski DA, Janisse DJ. Clinics in Sports Medicine 2004;23(1):157-67.

Functional bracing of the adult acquired flatfoot.

Marzano R. Clinics in Podiatric Medicine and Surgery 2007;24(4):645,56, vii.

De-stabilizing and training effects of foot orthoses in multiple sclerosis.

Ramdharry GM, et al. Multiple sclerosis 2006;12(2):219-26.

Effects of foot orthoses on patients with chronic ankle instability.

Richie DH. Journal of the American Podiatric Medical Association 2007;97(1):19-30.

Orthotic devices after stroke and other non-progressive brain lesions.

Tyson SF, Kent RM. Cochrane database of systematic reviews (Online) (1).1 2009: CD003694.

Orthotics and assistive devices in the treatment of upper and lower Limb osteoarthritis: An update.

Yonclas PP, et al. American Journal of Physical Medicine & Rehabilitation Suppl, 2006;85(11):S82-97.

Protheses

Comparison of interface pressures with pin and suction suspension systems.

Beil TL, Street GM. Journal of rehabilitation research and development 2004;41(6A):821-8.

Microprocessor prosthetic knees.

Berry D. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):91,113, vii.

Lower extremity socket design and suspension.

Carroll K. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):31-48.

Comparison of different microprocessor controlled knee joints on the energy consumption during walking in trans-femoral amputees: Intelligent knee prosthesis (IP) versus C-Leg.

Chin T, et al. Prosthetics and Orthotics International 2006;20(1): 73-80.

Effect of physical fitness on prosthetic ambulation in elderly amputees.

Chin T, Sawamura S, Shiba R. American Journal of Physical Medicine & Rehabilitation 2006;85(12):992-6.

Review of research on prosthetic devices for lower extremity amputation.

Collins DM, et al. Critical Reviews in Biomedical Engineering 2006;34(5): 379-438.

Componentry for lower extremity prostheses.

Friel K. Journal of the American Academy of Orthopaedic Surgeons 2005;13(5):326-35.

Review of secondary physical conditions associated with lower-limb amputation and long-term prosthesis use.

Gailey R, et al. Journal of Rehabilitation Research and Development 2008;45(1):15-29.

A comparative study of conventional and energy-storing prosthetic feet in high-functioning transfemoral amputees.

Graham LE, et al. Archives of Physical Medicine and Rehabilitation 2007;88(6):801-6.

A comparative study of oxygen consumption for conventional and energy-storing prosthetic feet in transfemoral amputees.

Graham LE, et al. Clinical Rehabilitation 2008;22(10-11):896-901.

Evaluation of function, performance, and preference as transfemoral amputees transition from mechanical to microprocessor control of the prosthetic knee.

Hafner B J, et al. Archives of Physical Medicine and Rehabilitation 2007;88(2):207-17.

Osseointegrated trans-femoral amputation prostheses: prospective results of general and condition-specific quality of life in 18 patients at 2-Year follow-up.

Hagberg K, et al. Prosthetics and Orthotics International 2008;32(1):29-41.

The effects of prosthetic foot design on physiologic measurements, self-selected walking velocity, and physical activity in people with transtibial amputation.

Hsu MJ, et al. Archives of Physical Medicine and Rehabilitation 2006;87(1):123-9.

Comparison of non-microprocessor knee mechanism versus C-Leg on prosthesis evaluation questionnaire, stumbles, falls, walking tests, stair descent, and knee preference.

Journal of rehabilitation research and development 2008;45(1):1-14.

Gait and balance of transfemoral amputees using passive mechanical and microprocessor-controlled prosthetic knees.

Kaufman KR, et al. Gait & Posture 2007;26(4):489-93.

Energy expenditure and activity of transfemoral amputees using mechanical and microprocessor-controlled prosthetic knees.

Kaufman KR, et al. Archives of Physical Medicine and Rehabilitation 2008;89(7):1380-5.

Targeted reinnervation for improved prosthetic function.

Kuiken T. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):1-13.

Clinical factors associated with prescription of a prosthetic limb in elderly veterans.

Kurichi JE, et al. Journal of the American Geriatrics Society 2007;55(6):900-6.

Progressive upper limb prosthetics.

Lake C, Dodson R. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):49-72.

Control of a six degree of freedom prosthetic arm after targeted muscle reinnervation surgery.

Miller LA, et al. Archives of Physical Medicine and Rehabilitation 2008;89(11): 2057-65.

Limb deficiency and prosthetic management. Decision making in prosthetic prescription and management.

Nelson VS, et al. Archives of Physical Medicine and Rehabilitation 2006;87(3):19, Suppl 1, S3-9.

A randomized controlled trial comparing functional outcome and cost efficiency of a total surface-bearing socket versus a conventional patellar tendon-bearing socket in transtibial amputees.

Selles RW, et al. Archives of Physical Medicine and Rehabilitation 2005;86(1):154,61; quiz 180.

Comparison between the C-Leg microprocessor-Controlled prosthetic knee and non-microprocessor control prosthetic knees: A preliminary study of energy expenditure, obstacle course performance, and quality of life survey.

Seymour R, et al. Prosthetics and Orthotics International 2007;31(1):51-61.

Pare and prosthetics: The early history of artificial limbs.

Thurston AJ. ANZ Journal of Surgery 2007;77(12):1114-9.

Changes in lower Extremity prosthetic practice.

Trower TA. Physical Medicine and Rehabilitation Clinics of North America 2006;17(1):23,30, v-vi.

Prosthetic feet: State-of-the-art review and the importance of mimicking human ankle-foot biomechanics.

Versluys R, et al. Disability and rehabilitation/Assistive technology 2009;4(2):65-75.

Prosthetic ankle-foot mechanism capable of automatic adaptation to the walking surface.

Williams RJ, Hansen AH, Gard SA. Journal of Biomechanical Engineering 2009;131(3):035002.

Prosthetic management of the partial foot amputee.

Yonclas PP, O'donnell CJ. Clinics in Podiatric Medicine and Surgery 2005;22(3):485-502.

Prosthetic feet: State-of-the-art review and the importance of mimicking human ankle-foot biomechanics.

Versluys R, Beyl P, Van Damme M, Desomer A, Van Ham R, Lefeber D. Disabil Rehabil Assist Technol Mar 2009;4(2):65-75.

Amputation-Related Issues

Clinical inquiries: What is the best way to manage phantom limb pain?

Black LM, Persons RK, Jamieson B. The Journal of family practice 2009;58(3):155-8.

Return to driving after lower-extremity amputation.

Boulias C, et al. Archives of Physical Medicine and Rehabilitation 2006;87(9):1183-8.

Return to work after lower limb amputation.

Burger H, Marincek C. Disability and Rehabilitation 2007;29(17):1323-9.

Partial hand amputation and work.

Burger H, Maver T, Marincek C. Disability and rehabilitation 2007;29(17):1317-21.

Self-delivered home-based mirror therapy for lower limb phantom pain.

American Journal of Physical Medicine & Rehabilitation 2009;88(1):78-81.

Reamputation, mortality, and health care costs among persons with dysvascular lower-limb amputations.

Dillingham TR, Pezzin LE, Shore AD. Archives of Physical Medicine and Rehabilitation 2005;86(3): 480-6.

Rehabilitation setting and associated mortality and medical stability among persons with amputations.

Dillingham TR, Pezzin LE. Archives of Physical Medicine and Rehabilitation 2008;89(6):1038-45.

Dermatologic conditions associated with use of a lower-extremity prosthesis.

Dudek NL, et al. Archives of Physical Medicine and Rehabilitation 2005;86(4):659-63.

Skin problems in an amputee clinic.

Dudek NL, Marks MB, Marshall SC. American Journal of Physical Medicine & Rehabilitation 2006;85(5):424-9.

The geriatric amputee.

Frieden RA. Physical Medicine and Rehabilitation Clinics of North America 2005;16(1):179-95.

Sexuality and amputation: A systematic literature review.

Geertzen JH, Van Es CG, Dijkstra PU. Disability and Rehabilitation 2009;31(7):522-7.

Postoperative management of lower extremity amputations.

Goldberg T. Physical Medicine and Rehabilitation Clinics of North America 17.1, 2006;17(1):173,80, vii.

Stability of phantom limb phenomena after upper limb amputation: A longitudinal study.

Hunter JP, Katz J, Davis KD. Neuroscience 2008;156(4):939-49.

Burns and amputations: A 24-year experience.

Kennedy PJ, et al. Journal of Burn Care & Research 2006;27(2):183-8.

The experience of phantom limb pain in patients with combat-related traumatic amputations.

Ketz AK. Archives of Physical Medicine and Rehabilitation 2008;89(6):1127-32.

Chronic low back pain in traumatic lower limb amputees.

Kulkarni J, et al. Clinical Rehabilitation 2005;19(1):81-6.

Survival rates in dysvascular lower limb amputees.

Kulkarni J, Pande S, Morris J. International Journal of Surgery 2006;4(4):217-21.

Long-term outcomes and survival of patients with bilateral transtibial amputations after rehabilitation.

Mac Neill HL, et al. American Journal of Physical Medicine & Rehabilitation 2008;87(3): 189-96.

Efficacy of bracing the lower limbs and ambulation training in children with myelomeningocele.

Mazur JM, Kyle S. Developmental medicine and child neurology 2004;46(5): 352-6.

Skin problems in lower limb amputees: A systematic review.

Meulenbelt HE, et al. Disability and Rehabilitation 2006;28(10):603-8.

Determinants of skin problems of the stump in lower-limb amputees.

Meulenbelt HE, et al. Archives of Physical Medicine and Rehabilitation 2009;90(1):74-81.

Low-Back Pain in transfemoral amputees: Is there a correlation with static or dynamic leg-length discrepancy?

Morgenroth DC, et al. American Journal of Physical Medicine & Rehabilitation 2009;88(2):108-13.

Advances in amputee care.

Pasquina PF, et al. Archives of Physical Medicine and Rehabilitation 2006;87(3):Suppl 1 2006: S34,43; quiz S44-5.

Falls sustained during inpatient rehabilitation after lower Limb amputation: prevalence and predictors.

Pauley T, Devlin M, Heslin K. American Journal of Physical Medicine & Rehabilitation 2006;85(6):521-32.

Heterotopic ossification following traumatic and combat-related amputations. Prevalence, risk factors, and preliminary results of excision.

Potter BK, et al. Journal of Bone and Joint Surgery 2007;89(3):476-86.

Identifying clinically meaningful improvement in rehabilitation of lower-limb amputees.

Resnik L. Medicine and health 2007;90(1):15-7.

Painful and nonpainful phantom and stump sensations in acute traumatic amputees.

Schley MT, et al. Journal of Trauma 2008;65(4):858-64.

Fluid collections in amputee stumps: A common phenomenon.

Singh R, Hunter J, Philip A. Archives of Physical Medicine and Rehabilitation 2007;88(5):661-3.

Managing the upper extremity amputee: A protocol for success.

Smurr LM, et al. Journal of Hand Therapy 2008;21(2):160,75; quiz 176.

The effectiveness of inpatient rehabilitation in the acute postoperative phase of care after transtibial or transfemoral amputation: Study of an integrated health care delivery system.

Stineman MG, et al. Archives of Physical Medicine and Rehabilitation 2008;89(10):1863-72.

The prevalence of osteoarthritis of the intact hip and knee among traumatic leg amputees.

Struyf PA, et al. Archives of Physical Medicine and Rehabilitation 2009;90(3):440-6.

Comparison of quality of life and functionality in patients with traumatic unilateral below knee amputation and salvage surgery.

Tekin L, et al. Prosthetics and Orthotics International 2009;33(1):17-24.

Early Treatment of trans-tibial amputees: Retrospective analysis of early fitting and elastic bandaging.

vanVelzen AD, et al. Prosthetics and Orthotics International 2005;29(1):3-12.

Quality of life of persons with lower-limb amputation during rehabilitation and at 3-month follow-Up.

Zidarov D, Swaine B, Gauthier-Gagnon C. Archives of Physical Medicine and Rehabilitation 2009;90(4):634-45.

Lower-extremity amputations in patients with diabetes: pre- and post-surgical decisions related to successful rehabilitation.

Davis BL, Kuznicki J, Praveen SS, Sferra JJ. Diabetes Metab Res Rev. May-Jun 2004;20Suppl 1:S45-50.